

Venous thromboembolism in pregnancy among Chinese women in a regional hospital in Hong Kong: a 20-year cross-sectional case-control study

Kwok Ting Jennifer LI, MBChB, MRCOG

Linus Lin Tai LEE, MBBS, MRCOG, FHKAM (Obstetrics and Gynaecology)

Lin Wai CHAN, MBChB, FHKAM (Obstetrics and Gynaecology), CertHKCOG (MFM)

Department of Obstetrics and Gynaecology, Pamela Youde Nethersole Eastern Hospital, Hong Kong SAR, China

Objective: To determine the prevalence and risk factors of venous thromboembolism (VTE) in pregnancy among Chinese women in a regional hospital.

Methods: Medical records of ethnic Chinese residents who delivered at ≥ 24 weeks of gestation or earlier with active resuscitation between January 2004 and December 2023 at Pamela Youde Nethersole Eastern Hospital were retrospectively reviewed. To determine VTE prevalence, cases occurring during pregnancy and the postpartum period (up to 6 weeks after delivery) were identified. Cases were excluded if associated with prophylactic anticoagulant use, negative imaging findings, or a history of VTE without active disease during pregnancy. To determine risk factors for VTE, cases were compared with matched controls (within 3 days of the delivery date) at a 1:5 ratio.

Results: In total, 22 cases of VTE occurring in the antepartum ($n=15$) or postpartum ($n=7$) period were identified. The prevalence was 0.5 per 1000 deliveries. Of the 22 cases, 16 were deep vein thrombosis, three were pulmonary embolism, and three involved both conditions. The most common symptom was limb swelling or pain ($n=17$). Two patients experienced cardiac arrest and underwent perimortem Caesarean section and subsequent mechanical thrombectomy by interventional radiologists. The remaining 20 patients with VTE received low-molecular-weight heparin treatment. The case and control groups were comparable, except that the case group had higher rates of substance use (13.6% vs 0.9%, $p=0.001$), varicose veins (4.5% vs 0%, $p=0.025$), preterm delivery (27.3% vs 7.3%, $p=0.005$), Caesarean section (63.6% vs 36.4%, $p=0.018$), blood transfusion (22.7% vs 2.7%, $p<0.001$), and stillbirth or neonatal death (4.5% vs 0%, $p=0.025$).

Conclusion: The prevalence of VTE in pregnancy among Chinese women at a regional hospital in Hong Kong was 0.5 per 1000 deliveries, lower than that reported internationally. The risks and benefits of pharmaco-prophylaxis should be weighed on an individual basis.

Keywords: East Asian people; Pregnancy complications, cardiovascular; Venous thromboembolism

Introduction

Pregnancy is a hypercoagulable state that develops as a protective mechanism to reduce haemorrhage at childbirth. It increases the risk of venous thromboembolism (VTE), which is a leading cause of maternal death¹. Prophylactic measures include compression stockings, early mobilisation, and pharmaco-prophylaxis; however, the latter is associated with bleeding risks. Risk-benefit analysis depends on the presumed prevalence of bleeding risks in different populations. The prevalences of VTE in pregnancy have been reported to be 1.4 per 1000 worldwide and 1.3 per 1000 in the Chinese population^{2,3}. In Hong Kong, the reported prevalence ranges from 0.4 to 1.88 per 1000 in tertiary units^{4,5}. We aimed to determine the prevalence and risk factors of VTE in pregnancy among Chinese women in a regional hospital across 20 years.

delivered at ≥ 24 weeks of gestation or earlier with active resuscitation between January 2004 and December 2023 at Pamela Youde Nethersole Eastern Hospital were retrospectively reviewed. Non-local residents were excluded, as were women with ectopic pregnancies, miscarriages, or abortions before fetal viability. During the study period, no standardised protocols or routine pharmaco-prophylaxis for VTE prevention in pregnancy were implemented. Clinicians provided prophylactic measures on an individual basis; non-pharmacological measures, such as compression stockings, were commonly prescribed for prolonged hospitalisation.

Data retrieved included anticoagulant use, limb ultrasound scans, computed tomography thoracic scans,

Methods

Medical records of ethnic Chinese residents who

Correspondence to: Dr Linus Lin Tai LEE

Email: leelintai@gmail.com

ventilation-perfusion scans, and diagnoses of VTE and related conditions between 280 days (40 weeks) before delivery and 100 days (~14 weeks) after delivery. There were no missing data.

To determine VTE prevalence, cases occurring during pregnancy and the postpartum period (up to 6 weeks after delivery⁶) were identified. Cases were excluded if associated with prophylactic anticoagulant use, negative imaging findings, or a history of VTE without active disease during pregnancy.

To determine risk factors for VTE, cases were compared with matched controls (within 3 days of the delivery date) at a 1:5 ratio to maximise statistical power⁷. The two groups were compared using the Chi-squared test for categorical variables and the *t* test for continuous variables. Statistical analysis was performed using SPSS (Windows version 29.0; IBM Corp, Armonk [NY], United States).

Results

Among 43 626 deliveries by Chinese residents, 22 cases of VTE occurring in the antepartum (*n*=15) or postpartum (*n*=7) period were identified. The prevalence was 0.5 per 1000 deliveries. Of the 22 cases, 16 were deep vein thrombosis, three were pulmonary embolism, and three involved both conditions (Table 1). The most common symptom was limb swelling or pain (*n*=17). Two patients experienced cardiac arrest and underwent perimortem Caesarean section and subsequent mechanical thrombectomy by interventional radiologists. The remaining 20 patients with VTE received low-molecular-weight heparin treatment. Only 11 of the 22 patients underwent thrombophilia screening after the VTE event.

The case and control groups were comparable, except that the case group had higher rates of substance use (13.6% vs 0.9%, *p*=0.001), varicose veins (4.5% vs 0%, *p*=0.025), preterm delivery (27.3% vs 7.3%, *p*=0.005), Caesarean section (63.6% vs 36.4%, *p*=0.018), blood transfusion (22.7% vs 2.7%, *p*<0.001), and stillbirth or neonatal death (4.5% vs 0%, *p*=0.025) [Table 2]. Indications for Caesarean section included placenta previa, maternal disease, twin pregnancy, failed induction after preterm prelabour rupture of membranes, and cardiac arrest.

Discussion

The incidence of VTE is lower among Asians, including ethnic Chinese^{8–10}. The prevalence of VTE in pregnancy in our hospital between 2004 and 2023 was 0.50

Table 1. Characteristics of the 22 cases of venous thromboembolism (VTE).

Characteristic	Value*
Occurrence	
Antepartum	15 (68.2)
>1st trimester	5 (33.3)
>2nd trimester	2 (13.3)
>3rd trimester	8 (53.3)
Postpartum	7 (31.8)
Presentation	
Limb swelling or pain	17 (77.3)
Cardiac arrest	2 (9.1)
Others (shortness of breath, stroke, tachycardia)	3 (13.6)
VTE type	
Deep vein thrombosis	16 (72.7)
Pulmonary embolism	3 (13.6)
Both	3 (13.6)
Site of deep vein thrombosis	<i>n</i> =19
Left lower limb	12 (63.2)
Right lower limb	5 (26.3)
Bilateral lower limbs	1 (5.3)
Left upper limb	1 (5.3)
Site of pulmonary embolism	<i>n</i> =6
Right main artery	1 (16.7)
Right lobar artery	2 (33.3)
Right segmental artery	1 (16.7)
Bilateral main arteries	1 (16.7)
Bilateral lobar arteries	1 (16.7)
Treatment	
Low-molecular-weight heparin	20 (90.9)
Thrombectomy	2 (9.1)
Thrombophilia screening	11 (50.0)

* Data are presented as No. (%) of cases

per 1000 deliveries (31.8% occurred postpartum), which was lower than the 1.3 per 1000 deliveries reported in the Chinese population³ and the 1.88 per 1000 deliveries (75% occurred postpartum) reported between 1998 and 2000 at Prince of Wales Hospital, Hong Kong⁴. It was similar to the 0.4 per 1000 deliveries (one case occurred postpartum) reported between 2004 and 2016 at Queen Mary Hospital, Hong Kong⁵. Regional differences within Hong Kong may reflect variations in socioeconomic status; higher socioeconomic status is associated with a decreased risk of VTE^{3,11}.

Table 2. Risk factors for venous thromboembolism (VTE).

Variable	VTE*		p Value
	Yes (n=22)	No (n=110)	
Age, y	33.5±4.3	33.2±4.4	-
Nulliparous	12 (54.5)	50 (45.5)	0.435
Parity			0.183
0	12 (54.5)	50 (45.5)	
1	7 (31.8)	49 (44.5)	
2	2 (9.1)	10 (9.1)	
3	0	0	
4	0	1 (0.9)	
≥5	1 (4.5)	0	-
Body mass index, kg/m ²	22.87±3.38	22.57±3.49	0.711
Smoker	4 (18.2)	8 (7.3)	0.104
Substance use	3 (13.6)	1 (0.9)	0.001
Comorbidities	1 (4.5)	0	0.025
Cardiac	0	1 (0.9)	0.653
Antiphospholipid syndrome	0	0	-
Systemic lupus erythematosus	0	1 (0.9)	0.653
History of VTE	0	0	-
Varicose veins	1 (4.5)	0	0.025
Others (sickle cell disease, urinary tract infection, irritable bowel disease)	0	0	-
Gestational diabetes	3 (13.6)	25 (22.7)	0.341
Pre-existing diabetes	1 (4.5)	1 (0.9)	0.202
Chronic hypertension	0	0	-
Pregnancy-induced hypertension	0	1 (0.9)	0.653
Pre-eclampsia	1 (4.5)	3 (2.7)	0.650
Intrauterine growth restriction	0	1 (0.9)	0.653
Artificial reproductive technology	2 (9.1)	5 (4.5)	0.385
Gestation at delivery, wk	38.1±2.1	38.9±1.5	0.056
Preterm delivery	6 (27.3)	8 (7.3)	0.005
Multiple pregnancies	2 (9.1)	3 (2.7)	0.153
Mode of delivery			0.048
Vaginal	6 (27.3)	60 (54.5)	
Instrumental	2 (9.1)	10 (9.1)	
Caesarean	14 (63.6)	40 (36.4)	0.018
Blood loss, mL	995±1974	311±210	0.119
Blood transfusion	5 (22.7)	3 (2.7)	<0.001
Manual removal of placenta	0	2 (1.8)	0.524
Stillbirth/neonatal death	1 (4.5)	0	0.025

* Data are presented as mean±standard deviation or No. (%) of cases

In the present study, inclusion of only local residents may have minimised cases with incomplete information or loss to follow-up. Most women remain in hospital for 3 to 5 days postpartum; most postpartum VTE occurs within the first week¹² and should therefore have been captured, particularly 90% of admissions are under Hospital Authority¹³. Although routine pharmaco-prophylaxis was not provided, mechanical measures (eg, compression stockings) for prolonged hospitalisations were offered on an individual basis. Routine pharmaco-prophylaxis during antepartum and postpartum periods has become more common in recent years¹⁴. Additionally, increased awareness of VTE, along with early mobilisation and adequate hydration postpartum, may reduce the incidence of postpartum VTE¹⁵. This trend of improved obstetric practice has also been observed internationally¹.

Although the bleeding risks associated with pharmaco-prophylaxis are not negligible^{14,16}, VTE in pregnancy remains uncommon in Hong Kong; thus, the risks and benefits of pharmaco-prophylaxis should be carefully weighed on an individual basis⁸.

Most VTEs were of deep vein thrombosis; symptoms predominantly affected the left lower limb, consistent with venous stasis in pregnancy¹⁷. Most cases occurred in the third trimester and postpartum periods. Maternal cardiac arrest is an uncommon presentation of pulmonary embolism in pregnancy¹⁸. Thrombectomy is preferred over thrombolysis, which may cause life-threatening haemorrhage¹⁹, highlighting the importance of the availability of interventional radiologists in obstetric units²⁰. Among patients with VTE, the prevalence of thrombophilia is high²¹. However, thrombophilia screening does not alter management and is therefore no longer advocated²².

Most patients with VTE had no known risk factors or comorbidities; only one had varicose veins. Risk factors for VTE include maternal age >35 years and obesity; lower prevalences of these risk factors may contribute to a reduced risk of VTE²³. Clinicians should maintain a high index of suspicion in pregnant women with symptoms suggestive of VTE, given that pregnancy itself is a risk factor^{22,24}. Substance abuse is associated with VTE in pregnancy, underscoring the importance of prevention and intervention efforts²⁵. Most postpartum VTEs are preceded by Caesarean section²³. Anticoagulation increases the risk of bleeding at delivery and postpartum, as well as the likelihood of blood transfusion, which itself is a risk factor for VTE²³.

The present study had several limitations. Considering the retrospective study design, patients' medical histories may not have been comprehensively recorded, and risk factors may have been underreported, potentially limiting the reliability of the results.

Conclusion

The prevalence of VTE in pregnancy among Chinese women at a regional hospital in Hong Kong was 0.5 per 1000 deliveries, lower than that reported internationally. The risks and benefits of pharmaco-prophylaxis should be weighed on an individual basis.

Contributors

All authors designed the study, acquired the data, analysed the data, drafted the manuscript, and critically revised the manuscript for important intellectual content. All authors had full access to the data, contributed to the study, approved the final version for publication, and take responsibility for its accuracy and integrity.

References

- Didembourg M, Morimont L, De Gottal E, Douchfils J. The maternal hemostatic shift: Understanding VTE risk in pregnancy and postpartum. *Thromb Res* 2026;257:109561. [Crossref](#)
- Kourlaba G, Relakis J, Kontodimas S, Holm MV, Maniadakis N. A systematic review and meta-analysis of the epidemiology and burden of venous thromboembolism among pregnant women. *Int J Gynaecol Obstet* 2016;132:4-10. [Crossref](#)
- Luo X, Shan D, Zhang L, et al. Incidence of maternal venous thromboembolism in China: a systematic review and meta-analysis. *Int J Gynaecol Obstet* 2023;163:75-88. [Crossref](#)
- Chan LY, Tam WH, Lau TK. Venous thromboembolism in pregnant Chinese women. *Obstet Gynecol* 2001;98:471-5. [Crossref](#)
- Huang D, Wong E, Zuo ML, et al. Risk of venous thromboembolism in Chinese pregnant women: Hong Kong venous thromboembolism study. *Blood Res* 2019;54:175-80. [Crossref](#)
- Kumarasinghe M, Herath MP, Hills AP, Ahuja KDK. Postpartum versus postnatal period: do the name and duration matter? *PLoS One* 2024;19:e0300118. [Crossref](#)
- Iwagami M, Shinozaki T. Introduction to matching in case-control and cohort studies. *Ann Clin Epidemiol* 2022;4:33-40. [Crossref](#)
- Liew NC, Alemany GV, Angchaisuksiri P, et al. Asian venous thromboembolism guidelines: updated recommendations for the prevention of venous thromboembolism. *Int Angiol* 2017;36:1-20. [Crossref](#)
- de Swiet M. Thromboembolism. *Clin Haematol* 1985;14:643-60. [Crossref](#)
- Tso SC, Wong V, Chan V, Chan TK, Ma HK, Todd D. Deep vein thrombosis and changes in coagulation and fibrinolysis after gynaecological operations in Chinese: the effect of oral contraceptives and malignant disease. *Br J Haematol* 1980;46:603-12. [Crossref](#)
- Jorgensen H, Horvath-Puho E, Laugesen K, Braekkan S, Hansen JB, Sorensen HT. Socioeconomic status and risk of incident venous thromboembolism. *J Thromb Haemost* 2021;19:3051-61. [Crossref](#)
- Tepper NK, Boulet SL, Whiteman MK, et al. Postpartum venous thromboembolism: incidence and risk factors. *Obstet Gynecol* 2014;123:987-96. [Crossref](#)
- Wong EL, Cheung AW, Leung MC, et al. Unplanned readmission rates, length of hospital stay, mortality, and medical costs of ten common medical conditions: a retrospective analysis of Hong Kong hospital data. *BMC Health Serv Res* 2011;11:149. [Crossref](#)
- Burd J, Zofkie A. Inpatient pharmacological thromboprophylaxis in the antepartum period: an argument for universal thromboprophylaxis. *Am J Obstet Gynecol MFM* 2025;7:101566. [Crossref](#)
- McLintock C, Brighton T, Chuniyal S, et al. Recommendations for the prevention of pregnancy-associated venous thromboembolism. *Aust N Z J Obstet Gynaecol* 2012;52:3-13. [Crossref](#)
- Schapkaitz E, Libhaber E, Gerber A, et al. A longitudinal study of thrombosis and bleeding outcomes with thromboprophylaxis in pregnant women at intermediate and high risk of VTE. *Clin*

Conflicts of interest

All authors have disclosed no conflicts of interest.

Funding/support

This study received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Data availability

All data generated or analysed during the present study are available from the corresponding author on reasonable request.

Ethics approval

This study was approved by the Hospital Authority Central Institutional Review Board (reference: CIRB-2024-601-5). The patients were treated in accordance with the tenets of the Declaration of Helsinki. The requirement for patient consent was waived by the Committee due to the retrospective nature of the research.

- Appl Thromb Hemost 2023;29:10760296231160748. [Crossref](#)
17. Chan WS, Spencer FA, Ginsberg JS. Anatomic distribution of deep vein thrombosis in pregnancy. *CMAJ* 2010;182:657-60. [Crossref](#)
 18. Martillotti G, Boehlen F, Robert-Ebadi H, Jastrow N, Righini M, Blondon M. Treatment options for severe pulmonary embolism during pregnancy and the postpartum period: a systematic review. *J Thromb Haemost* 2017;15:1942-50. [Crossref](#)
 19. Qadri S, Bilagi A, Sinha A, Connolly D, Murrin R, Bakour S. Acute management of massive pulmonary embolism in pregnancy. *Front Glob Womens Health* 2024;5:1473405. [Crossref](#)
 20. Baldwin HJ, Randall DA, Maher R, et al. Interventional radiology in obstetric patients: a population-based record linkage study of use and outcomes. *Acta Obstet Gynecol Scand* 2023;102:370-7. [Crossref](#)
 21. Tam WH, Ng MH, Yiu AK, Lau KM, Cheng GY, Li CY. Thrombophilia among Chinese women with venous thromboembolism during pregnancy. *Gynecol Obstet Invest* 2012;73:183-8. [Crossref](#)
 22. Ashraf N, Visweshwar N, Jaglal M, Sokol L, Laber D. Evolving paradigm in thrombophilia screening. *Blood Coagul Fibrinolysis* 2019;30:249-52. [Crossref](#)
 23. Alsheef MA, Alabbad AM, Albassam RA, et al. Pregnancy and venous thromboembolism: risk factors, trends, management, and mortality. *Biomed Res Int* 2020;2020:4071892. [Crossref](#)
 24. White RH. The epidemiology of venous thromboembolism. *Circulation* 2003;107(Suppl 1):I4-8. [Crossref](#)
 25. Bolnick JM, Rayburn WF. Substance use disorders in women: special considerations during pregnancy. *Obstet Gynecol Clin North Am* 2003;30:545-58. [Crossref](#)